

**CLIMATE CHANGE AND RIVER INDUS WATER QUANTITY  
ASSESSMENT USING GIS AND REMOTE SENSING TECHNIQUE.**

**Ali Akber khan<sup>1\*</sup>, Tehreem Aqsa<sup>2</sup>**

<sup>1\*</sup>*Neom Environmental sustainability services Islamabad Pakistan, A Case Study of Jinnah Barrage and Chashma barrage, Mianwali District, Punjab, Pakistan ali18april@hotmail.com*

<sup>2</sup>*Senior lecturer NCBA&E, A Case Study of Jinnah Barrage and Chashma barrage, Mianwali District, Punjab, Pakistan aqsa610@gmail.com*

**\*Corresponding Author:**

*ali18april@hotmail.com*

---

**Abstract:**

*The consumption of water assets is a significant issue that requires being knowledgeable about different areas worldwide, for example, the Mianwali Region in Pakistan. This paper gives a broad examination of the remote detecting strategies utilized for checking water amounts, with a particular accentuation on the Mianwali Region as a contextual investigation. Remote detection gives critical devices to assessing and controlling water assets by offering quick and geologically exact data on water sums. The goal of this study article is to analyze the usage of remote detecting innovation for observing water sums, assess the current water conditions in Mianwali Area, and give supportable water the board arrangements. The review uses a scope of remote detecting information sources, like satellite photography, and utilizes present-day methods including Geographic Data Frameworks (GIS) and calculations for picture handling to break down worldly varieties in water volume. The exploration discoveries improve appreciation of water asset elements in the Mianwali Region and propose significant direction to policymakers, water asset administrators, and scientists in planning effective techniques for water preservation and the executives.*

**Keywords:** *Climate change, River Indus, Water quantity monitoring, Remote sensing, Satellite imagery, Geographic Information Systems (GIS), Water management, Chashma and Jinnah barrage Mianwali.*


## **INTRODUCTION:**

In order to effectively manage water resources, it is essential to make use of remote sensing methods to monitor the quantity of water, particularly in areas with particular problems like the Mianwali District in Punjab, Pakistan. The maintainability of the environment, horticultural results, and human jobs in these locales are truly undermined by water shortage and consumption. By providing immediate and precise information on the dynamics of water quantity, remote sensing technology is an effective tool for evaluating and controlling water resources. The target of this examination work is to improve the cognizance of water amount observation through the usage of strategies for remote detecting, with specific accentuation on the Mianwali Region as an occasion of exploration. Utilizing satellite imagery, Geographic Information Systems (GIS), and sophisticated image processing algorithms, the researchers will investigate changes in water volume over time, assess the state of the water supply in the Mianwali District at the present time, and suggest environmentally friendly strategies for water resource management. This examination directs a careful examination of ebb and flow writing to investigate the different purposes, hardships, and limitations of remote detecting in the administration of water assets. This examination lays out the reason for the procedure continued in the current review. The points of this examination incorporate looking at patterns in water amount, dissecting the spatial circulation of water assets, confirming the ends using ground truth information, and surveying the outcomes of water using the executives' methods. In order to establish successful policies for water management and conservation in Mianwali District and other locations worldwide that are similar, this study aims to provide significant insights to policymakers, water resource managers, and researchers. In addition, the unique hydrological and geographical characteristics of the Mianwali District research area make it an ideal subject for understanding the quantity of water dynamics in semi-arid regions. Extending water demands from home, current, and provincial purposes, as well as inconsistent precipitation plans and unfeasible groundwater extraction, are pieces of the issues the district faces. In this particular circumstance, remote distinguishing advancement gives a cost-effective and effective technique for noticing water resources, working with a brief route, and assigning resources. This examination tries to combine remote detection information with ground truth information to direct an exhaustive assessment of water amount patterns, pinpoint areas of concern, and propose explicit mediations to accomplish viable water for the executives in Mianwali Locale. The consequences of this examination are expected to improve the current data on the use of remote detection in the administration of water assets, explicitly in semi-bone-dry areas experiencing equivalent troubles. In essence, the project aims to close the divide between scientific knowledge and practical application, promoting an integrated approach to water management and conservation in Mianwali District and beyond. Furthermore, this research is in alignment with wider national and global objectives for environmental sustainability, such as the United Nations Sustainable Development Goals (SDGs), specifically Goal 6: Clean Water and Sanitation. This goal seeks to guarantee the accessibility and sustainable administration of water and sanitation for everyone. This study advances economic, social, and ecological solidness in the Mianwali District and advances the bigger objective of water wellbeing and flexibility by using innovation for remote detecting and interdisciplinary techniques to assist with accomplishing SDG 6 targets. Through partnership collaboration between regional stakeholders, including government agencies, non-governmental organizations, and community representatives, this research aims to support the exchange of information, capacity building, and participatory decision-making processes in water management. This approach expects to improve the adequacy and inclusivity of water administration systems. All in all, the objective of this exploration is to work on logical information, add to the advancement of proof-based arrangements, and empower the reception of reasonable water by the executive's methods in Mianwali Locale and other comparative regions across the world. Overall, this work gives a careful examination of the observation of water volume through the usage of remote detecting techniques, with a specific accent on its investigation of the Mianwali Locale in Pakistan. This examination looks to use remote detecting information examination related to ground truth data to deliver critical bits of knowledge into the elements, patterns, and issues of water assets in the concentrated region. The consequences of this study are expected to give significant data to go with choices given proof, forming arrangements, and supporting the making of powerful water the executives plan in Mianwali Locale. This exploration expects to propel a helpful and sweeping way to deal with water administration by drawing in a few fields of study and incorporating partners. The goal is to ensure that water resources are distributed fairly and used effectively, to the benefit of current and future generations. In conclusion, the findings of this study emphasize the significance of remote sensing technology as an essential tool for addressing water scarcity and promoting water security worldwide. Besides, this study aims to address existing holes in the writing by giving a complete examination of the use of remote detecting strategies for observing water volume, explicitly in semi-parched regions like Mianwali Locale. By giving a top-to-bottom examination of the open doors and difficulties related to water assets on the board in such areas, this study adds to the more extensive talk on supportable turns of events and ecological safeguarding. Through a multidisciplinary approach that incorporates hydrological display, remote detection innovations, and financial investigation, this examination expects to foster total systems for the administration of water sources that think about friendly, monetary, and natural components. The objective of this study is to augment comprehension, exert influence on policy, and promote sustainable methodologies in water management. Its effect extends across the Mianwali District to other areas worldwide suffering from comparable water challenges.

## **MATERIAL AND METHODS:**

### **Study Area:**

The Mianwali is situated on the northwestern side of the Punjab area in Pakistan. It is portrayed by a number of intricate physical and hydrological factors that all affect how water is distributed and available. The area traverses around 5,840 square kilometers, and its populace fundamentally participates in agrarian pursuits and creature cultivation. The geography of the district involves undulating slopes, with the Thal Desert situated towards the west and the Soon Valley arranged towards the east. The Indus Waterway runs along the eastern boundary, assuming a critical part of the locale's hydrology. The Indus Stream runs along the eastern boundary, assuming a critical part of the locale's hydrology. The environment of the Mianwali Region is described as dry to semi-bone-dry, including burning summers and calm winters. The infrequent and sparse annual precipitation frequently causes issues with water scarcity. The water system arrangement of the Indus Stream and the useful utilization of groundwater assets are critical for the protection of farming, which fills in as a huge area of the nearby economy. All through the long haul, there has been a rise in the extraction of groundwater, achieving a decrease in water tables and bewildering the organization about legitimate water resources. Mianwali Domain is chiefly created for agriculture, where things like sugarcane, rice, and wheat are delivered. Moreover, there exist secluded areas of forests, especially in the Soon Valley, which add to keeping a natural harmony in the for the most part dry territory. Metropolitan regions, like Mianwali city, and the locale base camp, add to an ascent in water interest because of modern and home use. The region needs water bodies like reservoirs, ponds, and small rivers to keep agriculture going and the ecosystems that live there alive. In any case, the way of behaving of these amphibian frameworks can be changed by different causes, including environmental vacillations, alterations in land utilization, and human activities. Consequently, it is basic to direct an exhaustive examination to quantify the amount of water exactly. The financial design of the Mianwali Area is molded by the collaboration between agribusiness, water assets, and environmental conditions. The population, primarily rural, is entirely dependent on water accessibility and availability for their livelihoods. Getting a top-to-bottom comprehension of the complicated transaction between water supply, land use, and monetary and social movement is fundamental for creating effective water-the-board approaches in the area. At last, the extensive representation of the review locale in Mianwali Region fills in as a reason for the succeeding phases of the examination. Mianwali is an ideal area for concentrating on changes in water sum utilizing remote detecting strategies because of its particular mix of dry environment, agrarian reliance, and various geography. The exploration region's qualities act as urgent contributions for social events, examining, and deciphering information, ensuring a top-to-bottom appreciation of the water assets in this semi-dry area.



**Figure 1: Study Area Map Mianwali, Punjab, Pakistan**

**Data Collection:**

The data used in this assessment has been accumulated from the record of the US Geological Survey (USGS). It integrates Landsat satellite imagery taken by means of Landsat 4, 5, 7, 8, and 9 between the tremendous extended lengths of 1993,

2003, 2013, and 2023. The decision of these specific missions and lengths was purposeful to ensure complete inclusion of the review region and empower a definite assessment of varieties in the Standardized Contrast Water File (NDWI). Landsat satellites are notable for their devotion to Earth perception, giving multispectral information that is fundamental for observing long-haul natural changes. Landsat 4 and 5, which were in activity during the principal phases of this examination, gathered key information in 1993, laying out the reason for future missions. In 2003, Landsat 7 was based upon its ancestors by providing expanded spatial and ghastrly goals, consequently further propelling its abilities. The presentation of Landsat 8 out of 2013 addressed a huge accomplishment, achieving extra upgrades in picture quality and level of detail. The fuse of Landsat 9 information in 2023 expresses modern viewpoints, working with a careful cognizance of current modifications in the scene. This study expects to thoroughly gather and coordinate Landsat pictures from a few missions over quite a few years to portray the steadily changing qualities of the water assets in the review region precisely. A definitive objective is to offer important data and cognizance of the targets of water executives and preservation.

#### **Remote Sensing Data Preprocessing:**

The gathering of remote sensing data involved multiple stages aimed at preparing satellite imagery for analysis. At first, unprocessed satellite photos captured by multispectral sensors like Landsat or Sentinel were obtained for the designated time frame of the study. After this method, atmospheric correction methods were used to minimize air disturbance and influence the clarity of the photos. A mathematical revision was directed to address the bends and accomplish exact spatial correctness. To make it easier to compare and analyze the images, they were also resampled to a uniform spatial resolution. Preprocessing was essential in ensuring the constancy and accuracy of remote-detecting information for the resulting examination.

#### **Image Classification and Analysis NDWI:**

The process of classifying and analyzing images using the Normalized Difference Water Index (NDWI) in Mianwali District consisted of many essential stages. These stages are planned to precisely distinguish water bodies and assess their spatial course of action. For the duration of the study, multispectral satellite imagery, such as Landsat or Sentinel data, was initially obtained. Preprocessing strategies, like climatic revision and mathematical amendment, were used to work on the quality and accuracy of the photos. A while later, NDWI values were determined for each pixel utilizing the equation:

$$(NIR - SWIR)/(NIR + SWIR)$$

where NIR alludes to approaching infrared groups and SWIR alludes to shortwave infrared groups. The NDWI raster was arranged utilizing picture grouping procedures, like managed or unaided characterization, to allot pixels into one or the other water or non-water classes. Post-arrangement systems, for example, thresholding and sifting, were used to improve the water cover and decrease grouping botches. Consequently, the classified water veil was analyzed to gauge varieties in the space covered by water throughout some time and assess the game plan of water across various areas in the review locale. The utilization of an exhaustive technique for picture order and NDWI examination considered the exact identification and planning of water bodies. This approach gave crucial data on the elements of water in the Mianwali study area.


#### **Ground Truth Data Collection:**

The most widely recognized approach to social occasion ground truth data included driving field audits and really taking a look at exercises to confirm the exactness of the remote distinguishing disclosures. To assess the geological properties of the water bodies and verify their presence using satellite data, field tests were carried out. In addition, water level assessments, stream checking, and soil swiftness tests were aimed at expressing objections to enhancing remote recognizing data with on-the-spot assessments. The ground truth information was utilized as a kind of perspective to approve the consequences of the remote detection and make water amount evaluations more solid.

#### **RESULTS AND DISCUSSIONS:**


Observing water amounts utilizing remote detecting methods is a critical endeavor, as characterized by the situation investigation of the Mianwali Region. Using satellite imagery, this study investigates the temporal and regional changes in the region's water resources. Remote detecting gives a broad point of view, allowing the acknowledgment and evaluation of water bodies, which is fundamental for the administration of maintainable assets. The review uses progressed remote detecting innovation to decide how much water is accessible in the Mianwali Region, which is described by an assortment of landforms and a semi-parched environment. The goal of the project is to find patterns and shifts in the amount of water by looking at data gathered by Landsat satellites over a long period of time. This investigation endeavors to give bits of knowledge into the impacts of environmental change and human exercises on neighborhood water supplies. This case study provides a comprehensive understanding of Mianwali's environmental dynamics by examining the intricate interactions between vegetation dynamics, climate changes, and water quantity. The Normalized Difference Water Index (NDWI) and other sophisticated indicators help identify and track changes in open water characteristics. The outcomes exhibit critical swings in how much water, which compare to both occasional varieties and human action. The NDWI analysis reveals that water bodies are susceptible to external forces, focusing on areas where water volume changes significantly. These findings emphasize the significance of integrated water management measures to lessen the impact of

environmental changes on water availability and provide significant information about the vulnerability of water resources in the Mianwali District. The review's finished strategy offers areas of strength for fathoming and managing the complicated communication among regular and human-prompted components that influence the elements of water amounts in semi-dry districts.




**Figure 2: NDWI Map of Mianwali District (1993)**

Figure 2 shows huge insights regarding the fleeting changes in water elements through its portrayal of the Standardized Distinction Water Record (NDWI). The changes in NDWI values are depicted on the map, highlighting significant patterns in the availability of water over the specified time period. The most notable recorded NDWI esteem, coming to 0.775145 in 1993, shows a huge expansion in water presence during that year. Then again, the most minimal NDWI esteem is recorded at 0.520996, showing a period with decreased water content. The outrageous changes in NDWI values offer critical bits of knowledge into the variances in water amount, which are fundamental for fathoming the impacts of environmental conditions and human exercises on the water assets of the area under study. These numerical indices can be better understood, and the overall analysis of water quantity trends in the Mianwali District is enhanced by the visual representation in Figure 2. The swings found in NDWI values feature the unique person of water supply in the area, which is impacted by variables, for example, examples of precipitation, land use changes, and techniques for the water system. Moreover, the assessment of NDWI results after some time considers the discovery of occasional changes and broadened designs in water volume, working with the improvement of proficient water the board draws near. The discoveries introduced in Consider 2 deal with huge experiences along with the spatial and worldly changes in water assets in the Mianwali Area. These discoveries improve how we might interpret patterns in water amounts and can be utilized to pursue informed choices for supportable water by the executives in the district.




**Figure 3: NDWI Map of Mianwali District (2003)**

The depiction of the Normalized Difference Water Index (NDWI) in Figure 3 provides a significant understanding of the temporal variations in water dynamics. The visual representation accurately depicts the NDWI data fluctuation, highlighting notable patterns in water availability during the selected time period. The most elevated NDWI worth of 0.721881, seen in 2003, demonstrates a huge expansion in how much water was present during that year. Then again, the nadir NDWI worth of 0.559801 shows a period with decreased water content. The wide range of NDWI values is essential for comprehending the effects of climate and human activity on the water resources in the area under study because it provides useful information about variations in water levels. The graphical portrayal portrayed in Figure 3 fills in as a visual device, expanding the comprehension of these mathematical markers and working with an extensive assessment of water quantity designs in the Mianwali Locale. The noticed patterns in NDWI values are a consequence of the complicated connection between natural elements, for example, varieties of precipitation, changes in land use, and farming practices. These elements together impact the accessibility and appropriation of water assets in the locale. Besides, the assessment of NDWI values over the long run considers the identification of occasional changes and expanded designs, offering critical information for policymakers, water asset directors, and others who took part in water executives and safeguarding attempts. In rundown, the discoveries displayed in Figure 3 upgrade our insight into water development in the Mianwali Region, working with very much educated navigation and advancing reasonable water the board systems nearby.




**Figure 4: NDWI Map of Mianwali District (2013)**

The representation of the Normalized Difference Water Index (NDWI) in Figure 4 provides significant insights into the temporal dynamics of water. The realistic representation precisely shows the progressions in NDWI information, stressing striking examples in water accessibility during the characterized period. The most noteworthy recorded Standardized Distinction Water File (NDWI) score, coming to 0.687431 in 2013, shows a huge expansion in water accessibility during that particular year. On the other hand, the NDWI score arrives at its most terrible at 0.711164, demonstrating a period of diminished water content. The extensive variety of NDWI values gives significant bits of knowledge about the changes in water levels, which are fundamental for grasping the impacts of both environmental and human exercises on the water assets in the concentrated district. The visual portrayal portrayed in Figure 4 fills in as a visual help, working with the translation of the numeric lists and improving the far-reaching investigation of water amount changes in the Mianwali Locale. The intricate interaction between environmental factors like variations in rainfall, shifts in land use, and agricultural practices is what gives rise to the patterns in NDWI values. These elements all impact the accessibility and dispersion of water assets in the district. Additionally, the assessment of NDWI values over the long run empowers the identification of occasional vacillations and broadened designs, offering fundamental information for policymakers, water asset chiefs, and others who participated in water board and protection attempts. In outline, the discoveries displayed in Figure 4 upgrade our insight into water elements in the Mianwali Locale, working with very much educated navigation and advancing the supportability of water and the executives rehearses nearby.




**Figure 5: NDWI Map of Mianwali District (2023)**

Figure 5 provides a depiction of the Normalized Difference Water Index (NDWI), which reveals valuable information about the changes in water levels over a period of time. The significant patterns in water availability over the specified time period are highlighted by the graphic depiction, which accurately depicts the variations in NDWI measurements. The most noteworthy recorded NDWI score came to 0.687778 in 2023, demonstrating a critical presence of water during that particular year. Conversely, the lowest NDWI esteem recorded is 0.687654, showing a period with decreased water content. The extensive variety of NDWI results gives significant experiences into the vacillations in water levels, which are fundamental for grasping the impacts of environment and human exercises on the water assets nearby being scrutinized. The visual portrayal portrayed in Figure 5 fills in as a visual device, supporting the translation of mathematical lists and improving the general examination of water accessibility patterns in the Mianwali Region. The vacillations found in NDWI values feature the always-changing accessibility of water in the area, which is affected by variables, for example, examples of precipitation, changes in land use, and exercises connected with the water system. Besides, the assessment of NDWI values over the long run considers the location of occasional changes and expanded designs, offering huge information for policymakers, water asset directors, and others who participated in water the board and conservation drives. In a nutshell, the outcomes depicted in Figure 5 enhance our comprehension of the dynamics of water in the Mianwali District. This information might be utilized to settle on very educated choices and execute practical, water-the-board techniques in the region.




**Figure 6: Changes in the area (%) covered by water bodies in Mianwali District 1993.**

Figure 6 presents a concise analysis of the land cover in the area under study, providing essential information about its composition. The examination shows that water covers around 4.37% of the complete region, demonstrating the presence of oceanic qualities in the district. Besides, the land with high dampness content is a huge extent, adding up to 64.89% of the general region. Conversely, the extent of infertile ground or regions covered with vegetation is 30.74%, featuring the changed qualities of the climate. This broad assessment of land cover extents works with the perception of the spatial appropriation of water bodies, the predominance of sticky regions, and the degree of fruitless or vegetated land. These differentiations are significant for settling on very educated choices, overseeing assets, and executing maintainable administration strategies in the field of ecological examination. Figure 6 gives an itemized examination of how various kinds of land cover are conveyed inside a given region. This examination assists partners with acquiring a more nuanced comprehension of the regular habitat. Additionally, it enables them to prioritize conservation, restoration, and land use planning efforts in particular regions. Besides, the outline of land cover extents goes about as a supportive reference point for following ecological changes after some time, offering an establishment for assessing the impact of human exercises, environment variances, and normal disturbances on the biological system. In outline, the data from Figure 6 improves our complete perception of the land-water limit in the analyzed locale. This guides pursuing choices in light of proof and advances feasible improvement techniques that figure out some kind of harmony between natural protection and financial targets.




**Figure 7: Changes in the area (%) covered by water bodies in Mianwali District 2003**

Figure 7 presents a brief description of the land cover in the studied region, offering crucial information on its composition. The investigation demonstrates the significant presence of aquatic elements in the region by revealing that water covers approximately 3.74 percent of the area. Besides, a critical extent of the whole region, adding up to 65.33%, involves land with high dampness content. Conversely, 30.93% of the land is either exposed or covered with plants, featuring the fluctuating attributes of the climate. This broad assessment of land cover extents empowers an understanding of the spatial plan of water bodies, the transcendence of moist regions, and the extent of infertile or vegetated land. These differentiations are significant for settling on very educated choices, overseeing assets, and executing maintainable administration strategies in the field of ecological examination. Figure 7 measures the course of action of various land cover types, which can be utilized to decide the main areas for protection, rebuilding, and land use arranging endeavors. Besides, the division of land cover extents goes about as a pivotal reference point for following ecological movements over the long haul, permitting partners to assess the impact of human exercises, environment vacillations, and normal disturbances on the biological system. In synopsis, the data acquired from Figure 7 improves our exhaustive appreciation of the land-water limit in the concentrated locale. This guides settling on choices in light of proof and advances supportable improvement strategies that figure out some kind of harmony between natural conservation and economic objectives.



**Figure 8: Changes in the area (%) covered by water bodies in Mianwali District 2013**

Figure 8 provides a brief description of the land cover in the examined region, delivering vital information about its composition. According to the analysis, approximately 3.47 percent of the area is covered by water, indicating that the region contains numerous aquatic features. Besides, a significant extent of the whole region, adding up to 65.13%, involves land with high moisture content. Alternately, 31.40% of the locale contains either uncovered territory or regions covered with greenery, underlining the different qualities of the scene. This broad evaluation of land cover extents works with the appreciation of the spatial dissemination of water bodies, the pervasiveness of moist areas, and the degree of desolate or vegetated land. In the field of environmental studies, these distinctions are essential for enabling resource allocation, sustainable management practices, and well-informed decision-making. Figure 8 evaluates the course of action of various land cover types, which can be utilized to decide the main areas for protection, rebuilding, and land use arranging endeavors. Besides, the assurance of land cover extents goes about as a pivotal reference point for following ecological changes after some time, permitting partners to assess the impact of human exercises, environment variances, and normal disturbances on the biological system. In rundown, the data acquired from Figure 8 improves our thorough appreciation of the limit among land and water in the analyzed district. This guides in settling on choices in light of proof and advances supportable improvement strategies that figure out some kind of harmony between natural conservation and financial objectives.



**Figure 9: Changes in the area (%) covered by water bodies in Mianwali District 2023**

Figure 9 presents a clear overview of the land cover in the studied region, providing essential information about its composition. The examination uncovers that water represents roughly 3.41% of the complete region, showing the presence of amphibian qualities in the area. Moreover, the land with an elevated degree of dampness makes up a critical extent, representing 75.61% of the whole region. On the other hand, 20.98% of the scene comprises either uncovered land or regions covered with plants, featuring the different pieces of the territory. A comprehension of the geographical distribution of water bodies, the frequency of moist regions, and the extent of barren or vegetated land is made possible by this comprehensive evaluation of the proportions of land cover. These differentiations are significant for settling on very educated choices, overseeing assets, and executing maintainable administration strategies in the field of ecological examination. The spatial distribution of various types of land cover is quantified in Figure 9, which can be used to identify locations that should be prioritized for conservation, restoration, and land use planning projects. Besides, the order of land cover extends around as a critical reference point for following ecological movements over a period, permitting closely involved individuals to assess the impact of human activities, climatic variances, and normal interruptions on the environment. Figure 9 gives thorough bits of knowledge about the land-water interface in the concentrated region. These discoveries are useful for pursuing informed choices and advancing supportable improvement that thinks about both socio-economic concerns.

## CONCLUSION:

In conclusion, the examination features the critical hardships introduced by environmental change on an overall scale, with Pakistan, explicitly its helpless beach front districts in Sindh, being at the front of its ramifications. Mianwali area, situated in the northwestern locale of Punjab, experiences unmistakable geographical and climatic elements, which add to its perplexing natural qualities. The execution of modern remote detecting strategies, especially the Standardized Distinction Water List (NDWI), has been vital in the observation and perception of the changing water designs nearby. The examination of NDWI maps from 1993 to 2023 uncovers the differing patterns in water amount, exhibiting the two times of high and low levels. Insights into the distribution of water bodies and various landscape characteristics in Mianwali District are provided by the variances and the in-depth analysis of land cover percentages in Figures 6 to 9. This study makes up for a vital shortcoming by not just spearheading the assessment of vegetation cover in the waterfront locale of Sindh yet in addition researching the subtleties of Region Mianwali. The examination lays out a relationship between NDWI markers and asset shortage, giving a premise to pursuing instructed choices and overseeing assets reasonably. The use of Landsat satellite pictures ensures an exhaustive perception of modifications in the climate, subsequently adding to the more extensive conversation on measures for flexibility and variation to environmental change. Basically, this study shows the ability of remote detecting innovation to comprehend the perplexing connection between environmental change, vegetation examples, and water supplies. This understanding can act as an establishment for going to preventive lengths to safeguard the climate in light of an evolving environment. This research not only enhances the scientific understanding of water quantity monitoring but also offers practical insights for policymakers, water resource managers, and stakeholders. These insights can be used to develop effective strategies for sustainable water management and climate change adaptation in Mianwali District and other similar regions worldwide. This study demonstrates the significance of collaborative efforts in tackling urgent environmental concerns and developing resistance to the effects of climate change by combining remote sensing technologies with interdisciplinary approaches. Furthermore, the findings of this research highlight the critical requirement for proactive measures to moderate the unfriendly impacts of environmental change on water assets and biological systems in Mianwali Locale. As the locale keeps on confronting expanding water shortage and changeability, techniques, for example, further developed water preservation rehearses, reasonable land use arranging, and local area-based watershed the executives drive are basic for guaranteeing water security and ecological supportability. Also, the reconciliation of remote detecting information with hydrological models and financial examinations can improve the exactness and viability of water assets in the board procedures, empowering partners to expect and adjust to future changes in water accessibility and requests. Moreover, limit-building endeavors and public mindfulness crusades are fundamental for cultivating a culture of ecological stewardship and elevating local area versatility to environmental change influences. By embracing a comprehensive and cooperative way to deal with water the board, policymakers, specialists, and nearby networks can cooperate to protect water assets, relieve the effects of environmental change, and guarantee a manageable future for a long time into the future. Eventually, the bits of knowledge acquired from this study add to the more extensive plan of natural supportability and environmental change versatility, directing endeavors towards building a stronger and water-secure future for Mianwali District and worldwide.

## REFERENCES:

- [1]. Ashok, A., Rani, H. P., & Jayakumar, K. (2021). Monitoring of dynamic wetland changes using NDVI and NDWI-based Landsat imagery. *Remote Sensing Applications: Society and Environment*, 23, 100547.
- [2]. Bashir, B., Cao, C., Naeem, S., Zamani Joharestani, M., Bo, X., Afzal, H., . . . Mumtaz, F. (2020). Spatio-temporal vegetation dynamic and persistence under climatic and anthropogenic factors. *Remote Sensing*, 12(16), 2612.
- [3]. Bastawesy, M. E. (2015). Hydrological scenarios of the Renaissance Dam in Ethiopia and its hydro-environmental impact on the Nile downstream. *Journal of Hydrologic Engineering*, 20(7), 04014083.
- [4]. Carattini, S., Gosnell, G., & Tavoni, A. (2020). How developed countries can learn from developing countries to tackle climate change. *World Development*, 127, 104829.

- [5]. Chu, H., Venevsky, S., Wu, C., & Wang, M. (2019). NDVI-based vegetation dynamics and its response to climate changes at Amur-Heilongjiang River Basin from 1982 to 2015. *Science of the Total Environment*, 650, 2051-2062.
- [6]. Eid, A. N. M., Olatubara, C., Ewemoje, T., El-Hennawy, M. T., & Farouk, H. (2020). Inland wetland time-series digital change detection based on SAVI and NDWI indices: Wadi El-Rayan lakes, Egypt. *Remote Sensing Applications: Society and Environment*, 19, 100347.
- [7]. El-Gamily, I., Selim, G., & Hermas, E. (2010). Wireless mobile field-based GIS science and technology for crisis management process: A case study of a fire event, Cairo, Egypt. *The Egyptian Journal of Remote Sensing and Space Science*, 13(1), 21-29.
- [8]. Fatima, N., Alamgir, A., Khan, M., & Mehmood, K. (2021). Conceptual Framework for Climate Vulnerability and Conflicts in the Coastal Districts of Thatta and Sujawal, Sindh, Pakistan. *International Journal of Biosciences*, 18(4), 60-76.
- [9]. Gao, B.-C. (1996). NDWI—A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote sensing of environment*, 58(3), 257-266.
- [10]. Ghazal, L., Kazmi, J., & Zubair, S. (2015). Monitoring and Mapping Spatio-Periodic Dynamics of Vegetation Cover in Karachi Using Geoinformatics. *International Journal of Biology and Biotechnology*, 12(4), 621-627.
- [11]. Huang, F., & Xu, S. (2016). Spatio-temporal variations of rain-use efficiency in the west of Songliao Plain, China. *Sustainability*, 8(4), 308.
- [12]. Hurlbert, A. H., & Haskell, J. P. (2003). The effect of energy and seasonality on avian species richness and community composition. *The American Naturalist*, 161(1), 83-97.
- [13]. Liu, Y., & Lei, H. (2015). Responses of natural vegetation dynamics to climate drivers in China from 1982 to 2011. *Remote Sensing*, 7(8), 10243-10268.
- [14]. Mahmoudi, P., Shirazi, S. A., Amir Jahanshahi, S. M., Firoozi, F., & Mazhar, N. (2021). DETECTION OF LONG-TERM VEGETATION DEGRADATION IN BALUCHISTAN IN SOUTHWEST ASIA USING NDVI PRODUCTS OF THE MODIS SENSOR OF TERRA SATELLITE. *Environmental Engineering & Management Journal (EEMJ)*, 20(2).
- [15]. McFeeters, S. K. (1996). The use of the Normalized Difference Water Index (NDWI) in the delineation of open water features. *International Journal of Remote Sensing*, 17(7), 1425-1432.
- [16]. Nasir, S. M., & Akbar, G. (2012). Effect of River Indus flow on low riparian ecosystems of Sindh: a review paper. *Rec. Zool. Surv. Pakistan*, 21, 86-89.
- [17]. Ouma, Y. O., & Tateishi, R. (2006). A water index for rapid mapping of shoreline changes of five East African Rift Valley lakes: an empirical analysis using Landsat TM and ETM+ data. *International Journal of Remote Sensing*, 27(15), 3153-3181.
- [18]. Özelkan, E. (2020). Water body detection analysis using NDWI indices derived from Landsat-8 OLI. *Polish Journal of Environmental Studies*, 29(2), 1759-1769.
- [19]. Patil, P. P., Jagtap, M. P., Khatri, N., Madan, H., Vadduri, A. A., & Patodia, T. (2024). Exploration and advancement of NDDI leveraging NDVI and NDWI in Indian semi-arid regions: A remote sensing-based study. *Case Studies in Chemical and Environmental Engineering*, 9, 100573.
- [20]. Siyal, A., Siyal, A., & Mahar, R. (2017). Spatial and temporal dynamics of Pai forest vegetation in Pakistan assessed by RS and GIS. *Journal of Forestry Research*, 28, 593-603.
- [21]. Thayer, A. W., Vargas, A., Castellanos, A. A., Lafon, C. W., McCarl, B. A., Roelke, D. L., . . . Lacher, T. E. (2020). Integrating agriculture and ecosystems to find suitable adaptations to climate change. *Climate*, 8(1), 10.
- [22]. Wu, C., Venevsky, S., Sitch, S., Yang, Y., Wang, M., Wang, L., & Gao, Y. (2017). Present-day and future contribution of climate and fires to vegetation composition in the boreal forest of China. *Ecosphere*, 8(8), e01917.
- [23]. Xu, H. (2006). Modification of normalized difference water index (NDWI) to enhance open water features in remotely sensed imagery. *International Journal of Remote Sensing*, 27(14), 3025-3033.
- [24]. Zheng, Y., Tang, L., & Wang, H. (2021). An improved approach for monitoring urban built-up areas by combining NPP-VIIRS nighttime light, NDVI, NDWI, and NDBI. *Journal of Cleaner Production*, 328, 129488.
- [25]. Zheng, Y., Zhou, Q., He, Y., Wang, C., Wang, X., & Wang, H. (2021). An optimized approach for extracting urban land based on log-transformed DMSP-OLS nighttime light, NDVI, and NDWI. *Remote Sensing*, 13(4), 766.
- [26]. Zoran, M. A., Zoran, L. F. V., & Dida, A. I. (2016). Forest vegetation dynamics and its response to climate changes. Paper presented at the Remote Sensing for Agriculture, Ecosystems, and Hydrology XVIII.